arXiv:cond-mat/0408169v1 [cond-mat.other] 8 Aug 2004

Low-temperature crossover in the momentum distribution of cold atomic gases in one dimension

Vadim V. Cheianov,[1] H. Smith,[2] and M. B. Zvonarev[2]
[1]*NORDITA, Blegdamsvej 17, DK-2100 Copenhagen, Denmark*
[2]*Ørsted Laboratory, H. C. Ørsted Institute, Universitetsparken 5, DK-2100 Copenhagen, Denmark*

The momentum distribution function for the two-component 1D gases of bosons and fermions is studied in the limit of strong interatomic repulsion. A pronounced reconstruction of the distribution is found at a temperature much smaller than the Fermi temperature. This new temperature scale, which equals the Fermi temperature divided by the dimensionless coupling strength, is a feature of the two-component model and does not exist in the one-component case. We estimate the parameters relevant for the experimental observation of the crossover effect.

I. INTRODUCTION

During the last few years the great advances in the art of cooling and trapping atomic gases have made it possible to study experimentally the behavior of fermion and boson gases in lower dimensions[1]. Of particular interest are the quasi-one-dimensional systems which can be realized as one-dimensional (1D) tubes in a two-dimensional optical lattice [2, 3, 4, 5]. The ground states of such systems are characterized by the ratio between the interaction energy and the kinetic energy. At high densities, the kinetic energy dominates and the system can be described by mean-field theory. At low densities, one enters a regime of strong coupling where the interaction energy dominates the kinetic energy. The difference between this strongly interacting regime and the regime of weak interactions is more than just quantitative. One should expect that strong interactions result in phenomena absent in the weakly interacting case. The question of what these phenomena are and their experimental consequences is essential for our understanding of low-dimensional systems and, more generally, the physics of strongly interacting matter.

It is evident that the presence of internal degrees of freedom enriches the phase diagram of a physical system and may even give rise to completely new phenomena. One-dimensional systems in general and one-dimensional quantum gases in particular, are no exceptions to this rule. In this paper we discuss a phenomenon in quantum gases which is specific to the strongly interacting regime and the presence of internal degrees of freedom. We show that, due to the emergence of a soft propagating mode in the limit of strong repulsion, one-dimensional gases (both bosonic and fermionic) undergo a dramatic reconstruction of the one-particle momentum distribution as the temperature is varied on a very small scale. This temperature scale is proportional to the propagation velocity of the soft mode and vanishes in the limit of infinitely strong interparticle repulsion.

The paper is organized as follows. In Section II we give a qualitative comparison of one- and two-component quantum gases in 1D and introduce the relevant energy scales. We show that in the two-component case there exists a new low-temperature regime where the behavior of the quantum gas should be qualitatively different from that at zero temperature. Sections III and IV are technical. In Section III we present an exactly solvable model, which grasps the physics discussed in Section II. We give an exact representation for the one-particle density matrix in this model. In Section IV we analyze short- and long-distance asymptotics of the density matrix and discuss the numerical recipe for evaluating it at intermediate distances. Section V contains the main result of the paper: The effect of the low temperature reconstruction of the momentum distribution function is demonstrated for both boson and fermion gases. In Section VI we discuss the possibility of observing this reconstruction effect experimentally and estimate the magnitude of the coupling strength for such an experiment in terms of the parameters characterizing highly elongated harmonic traps.

II. ONE-COMPONENT VERSUS TWO-COMPONENT GASES IN ONE DIMENSION

One-dimensional *uniform* gases with *zero-range* interaction serve as a test ground for a wide class of one-dimensional systems. This was realized long ago by Lieb and Liniger, who used the Bethe-ansatz method [6] to obtain the exact wave functions and spectrum of the one-component boson gas [7]. They considered the following Hamiltonian for N identical bosons of mass m and chemical potential μ,

$$H = -\frac{\hbar^2}{2m}\sum_{i=1}^{N}\frac{\partial^2}{\partial x_i^2} + g_{1\mathrm{D}}\sum_{1\le i<j\le N}\delta(x_i - x_j) - \mu N. \quad (1)$$

The interaction constant $g_{1\mathrm{D}}$ has the dimension of energy times length and is conventionally written in terms of a characteristic length $a_{1\mathrm{D}}$ according to

$$g_{1\mathrm{D}} = \frac{2\hbar^2}{m a_{1\mathrm{D}}}. \quad (2)$$

The dimensionless coupling strength γ involves the number of particles per unit length, $n_{1\mathrm{D}} = N/L$, where L is the length of the 1D system, and is given by

$$\gamma = \frac{m g_{1\mathrm{D}}}{\hbar^2 n_{1\mathrm{D}}} = \frac{2}{n_{1\mathrm{D}} a_{1\mathrm{D}}}. \quad (3)$$

Thus when the density decreases, the coupling strength increases, and one approaches the limit of impenetrable bosons, the so-called Tonks-Girardeau gas [8]. In this limit the thermodynamic properties of the system are the same as those of a one-dimensional gas of non-interacting one-component fermions with the Fermi energy

$$E_{\rm F} = \pi^2 \frac{\hbar^2 n_{\rm 1D}^2}{2m}. \tag{4}$$

Henceforth we set $\hbar = 2m = 1$. These constants will be restored in Section VI.

Similarly to the one-component case, one can consider a *two-component* gas of fermions (bosons). Such gases consist of particles with some internal quantum parameter σ, which can take two values. For brevity we refer to this parameter as "spin", and label the corresponding states by $\sigma =\uparrow, \downarrow$. In the situation where the interaction constant $g_{\rm 1D}$ is tuned to be independent of the "spin" index, we take the first-quantized Hamiltonian of the system to be the same as for the one-component case, Eq. (1). The information about the system being two-component is encoded in the symmetry of the coordinate wave function. To introduce the "spin" indices in the Hamiltonian explicitly one can recast it in the second quantized form using the particle creation and annihilation operators $\psi^\dagger_\sigma(x)$ and $\psi_\sigma(x)$. Written in the second-quantized form the interaction term in Eq. (1) is

$$\frac{g_{\rm 1D}}{2} \int dx\, n_\uparrow(x) n_\downarrow(x) \tag{5}$$

for two-component fermions, and

$$\frac{g_{\rm 1D}}{4} \int dx\ : n(x)^2 : \tag{6}$$

for two-component bosons. Here $n_\sigma(x) = \psi^\dagger_\sigma(x)\psi_\sigma(x)$ is the density of particles of type σ; the total density $n(x)$ is $n(x) = n_\uparrow(x) + n_\downarrow(x)$, and the symbol :: stands for the normal ordering. For interactions of this form the total number of particles N and the total number $N_{\uparrow(\downarrow)}$ of "spin" up (down) particles are good quantum numbers. Note, that the operator (5) is the most generic form of contact interaction for fermions, which is a consequence of the Pauli principle. In the bosonic case, however, the operator (6) represents a special situation where the three coupling constants are tuned to be equal.

The density $n_{\rm 1D}$ is controlled by the chemical potential μ, Eq. (1). In the infinite γ limit the relation between $n_{\rm 1D}$ and μ does not depend on the statistics of particles and has the following simple form

$$\pi n_{\rm 1D} = \sqrt{\mu}. \tag{7}$$

For the one-component gas this result can be found in *e.g.* Refs. [8], for the two-component gas in Ref. [9]. It follows from Eq. (7) that $\sqrt{\mu}$ has a dimension of inverse length. Hereafter we measure distances in units of $1/\sqrt{\mu}$.

What temperature scales characterize 1D gases with the Hamiltonian (1) in the limit of strong repulsion, $\gamma \gg 1$? For the one-component system there is only one temperature scale: the Fermi temperature $T_{\rm F} = E_{\rm F}/k_{\rm B}$, where $k_{\rm B}$ is the Boltzmann constant. Low temperatures $T \ll T_{\rm F}$ correspond to the degenerate quantum gas. The inclusion of "spin" gives rise to another characteristic temperature $T_0 \ll T_F$, defined by the bandwidth for the "spin" excitations. This bandwidth can be estimated as follows. Consider the sector of the Hilbert space with a given number of particles N and a polarization $M = N_\uparrow - N_\downarrow$. Denote by $E(M)$ the ground state energy of the Hamiltonian (1) in this sector. For fermions the function $E(M)$ has its absolute minimum at $M = 0$ [10] in agreement with the Lieb-Mattis theorem [11]. For bosons the function $E(M)$ reaches its absolute minimum at $|M| = N$ [12], giving rise to the demixing transition [13]. The energy difference per particle between the polarized (demixed) and unpolarized (mixed) phases

$$\epsilon = \frac{|E(N) - E(0)|}{N} \tag{8}$$

gives an estimate for the bandwidth of "spin" excitations: $T_0 \sim \epsilon/k_{\rm B}$. In the large γ limit ϵ is easily found from the thermodynamic Bethe ansatz [10]. For fermions one finds that to the leading order in $1/\gamma$

$$\epsilon = \frac{8 \ln 2}{3} \frac{E_{\rm F}}{\gamma}. \tag{9}$$

One can see that the characteristic temperature scale for "spin" excitations is given by $T_0 = T_{\rm F}/\gamma$. The same estimate is valid for bosons.

The two scales T_0 and $T_{\rm F}$ are well separated when γ is large enough. In this situation one can distinguish between two different quantum (which means $T \ll T_{\rm F}$) regimes: $T < T_0$ and $T > T_0$. The first regime has been studied extensively and is conventionally described by the Luttinger Liquid (LL) theory [14]; the results on the second one appeared very recently [15, 16, 17] for the limiting case $T_0 \ll T \ll T_{\rm F}$ and are qualitatively different from the LL theory predictions. This can be understood by noticing that for $T > T_0$ "spin" degrees of freedom are strongly "disordered" thus violating the LL theory applicability conditions. At the same time, the density degrees of freedom are not affected by the temperature until it becomes of the order of $T_{\rm F}$.

III. ONE-PARTICLE DENSITY MATRIX: AN EXACT REPRESENTATION

In this Section we investigate the one-particle density matrix of the two-component strongly repulsive ($\gamma \to +\infty$) 1D gases with the Hamiltonian (1) for the two limiting cases $T \ll T_0$ and $T_0 \ll T$ of the two quantum regimes discussed in Section II. Our considerations will be based on the exact solution of the model Hamiltonian (1) in the infinite γ limit. The two different regimes $T \ll T_0$ and $T_0 \ll T$ are accessed by choosing the appropriate order of limits $T \to 0$ and $\gamma \to \infty$, as described in

[15]: To ensure $T_0 \ll T$ one should take the limit $\gamma \to \infty$ first and then set $T = 0$ in the wave functions and spectra of the model, while for the LL regime $T \ll T_0$ one should take the limit $T \to 0$ first.

For two-component systems the one-particle density matrix is defined as

$$\rho(x) = \frac{1}{2}\left[\langle \psi_\uparrow^\dagger(x)\psi_\uparrow(0)\rangle + \langle \psi_\downarrow^\dagger(x)\psi_\downarrow(0)\rangle\right]. \tag{10}$$

Since the interaction terms Eqs. (5) and (6) are invariant with respect to "spin" rotation, one has $\langle \psi_\uparrow^\dagger(x)\psi_\uparrow(0)\rangle = \langle \psi_\downarrow^\dagger(x)\psi_\downarrow(0)\rangle$. The density of particles is $n_{1\mathrm{D}} = 2\rho(0)$. Both physical intuition and crucial technical advance come from the observation that in the infinite γ limit the exact Bethe-ansatz many-particle eigenfunctions of the Hamiltonian (1) factorize into a coordinate part which is similar to the wave function of non-interacting spinless fermions and a spin part which corresponds to an eigenstate of the isotropic Heisenberg chain. Using this factorization Ogata and Shiba [18] derived an explicit formula for the density matrix (10) in the case of two-component fermions on a lattice (the Hubbard model). The two-component fermion gas being the limit of the Hubbard Hamiltonian for the vanishing filling factor, Ogata and Shiba's formula can be used. It also has a straightforward extension to the bosonic case. Both bosonic and fermionic cases are discussed in the paragraph below.

Using the results of Ref. [18] we write the density matrix $\rho(x)$ as

$$\rho(x) = \frac{1}{2}\langle \chi^\dagger(x)\omega(\mathcal{N}_x)\chi(0)\rangle_{\mathcal{F}}, \tag{11}$$

where the average $\langle\rangle_{\mathcal{F}}$ is taken over the ground state of the system of non-interacting one-component fermions with creation and annihilation operators $\chi^\dagger$ and χ, respectively. The density of these fermions coincides with the density of physical particles $n_{1\mathrm{D}}$. The operator

$$\mathcal{N}_x = \int_0^x dx' \chi^\dagger(x')\chi(x') \tag{12}$$

counts the number of particles between points 0 and x. Assuming lattice regularization with an infinitesimal lattice constant, $\mathcal{N}_x$ is a Hermitian operator with integer eigenvalues. To define the function $\omega(N)$ consider an infinite isotropic Heisenberg chain. Its Hilbert space is spanned by vectors $|\ldots\sigma_1\sigma_2\ldots\sigma_j\ldots\rangle$, where $\sigma_j = \uparrow, \downarrow$ is the spin at the j-th site. In this space one may introduce a string operator Ω_N, which acts as a cyclic shift operator on the string of spins of length N:

$$\begin{aligned}\Omega_N|\ldots\sigma_j\sigma_{j+1}\ldots\sigma_{j+N-2}\sigma_{j+N-1}\sigma_{j+N}\ldots\rangle = \\ |\ldots\sigma_{j+N-1}\sigma_j\sigma_{j+1}\ldots\sigma_{j+N-2}\sigma_{j+N}\ldots\rangle.\end{aligned} \tag{13}$$

The function $\omega(N)$ is defined as follows

$$\omega(N) = e^{i(\pi-\phi)N}\langle\Omega_N\rangle_{\mathcal{H}}. \tag{14}$$

Here ϕ is a statistical angle: $\phi = 0$ for bosons and $\phi = \pi$ for fermions. In the case $T \ll T_0$ the average $\langle\rangle_{\mathcal{H}}$ in Eq. (14) is taken over the ground state of the isotropic Heisenberg chain, which is antiferromagnetic for the fermion gas and ferromagnetic for the boson gas. In the case $T \gg T_0$, the average $\langle\,\rangle_{\mathcal{H}}$ is to be taken over all possible spin configurations.

For the boson gas at exactly zero temperature and both fermion and boson gases at $T \gg T_0$ the calculation of $\omega(N)$ is relatively simple. For bosons at zero temperature the average $\langle\,\rangle_{\mathcal{H}}$ in Eq. (14) is taken over the fully polarized ferromagnetic ground state of the spin chain which gives

$$\omega(N) = e^{i\pi N}. \tag{15}$$

For bosons and fermions at $T \gg T_0$ the average $\langle\,\rangle_{\mathcal{H}}$ in Eq. (14) runs over all spin configurations. This gives

$$\omega(N) = e^{i(\pi-\phi)N}2^{-N}. \tag{16}$$

In both equations Eqs. (15) and (16) the function $\omega(N)$ is an exponential of N : $\omega(N) \sim e^{\lambda N}$, for which the average (11) can be expressed in terms of the Fredholm determinant. More precisely, one obtains the following representation for the density matrix (10):

$$\frac{\rho(x)}{\rho(0)} = \det(\hat{I} + \hat{V}_1 + \hat{V}_2)(x) - \det(\hat{I} + \hat{V}_1)(x). \tag{17}$$

Here the determinant

$$\begin{aligned}\det(\hat{I}+\hat{V}) = \sum_{N=0}^{\infty}\frac{1}{N!}\int_{-1}^{1}\mathrm{d}k_1\ldots\int_{-1}^{1}\mathrm{d}k_N \\ \times\det\begin{bmatrix} V(k_1,k_1) & \cdots & V(k_1,k_N) \\ \vdots & \ddots & \vdots \\ V(k_N,k_1) & \cdots & V(k_N,k_N)\end{bmatrix}\end{aligned} \tag{18}$$

is the Fredholm determinant of a linear integral operator $\hat{V}$ with the kernel defined on $[-1,1]\times[-1,1]$. In our case the kernels V_1 and V_2 are:

$$V_1(k,p) = \alpha\frac{\sin\frac{x}{2}(k-p)}{\pi(k-p)} \tag{19}$$

$$V_2(k,p) = \frac{1}{2}\exp\{-i\frac{x}{2}(k+p)\} \tag{20}$$

The values of the parameter α are given in the Table I. Recall that we measure distances in units of $1/\sqrt{\mu}$, Eq. (7)

It was mentioned above Eq. (15) that the ground state of the two-component bosons is fully polarized in "spin" [12]. Therefore, at $T \ll T_0$ the bosons become effectively one-component (compare first and second lines in the Table I) and we turn back to the original Lieb-Liniger model [7]. The determinant representation (17) in this model was obtained by Lenard [19]. For the two-component fermion gas at $T \gg T_0$ Eq. (17) was obtained in Ref. [20]. Izergin and Pronko calculated both bosonic and fermionic density matrices in Ref. [9].

	α
one-component boson gas, $T = 0$	-2
two-component boson gas, $T \ll T_0$	-2
two-component boson gas, $T \gg T_0$	$-3/2$
two-component fermion gas, $T \ll T_0$	—
two-component fermion gas, $T \gg T_0$	$-1/2$

Table I: The values of the parameter α entering Eq. (19) are given. For the two-component fermion gas at $T \ll T_0$ no exact results on $\rho(x)$ are known.

IV. ONE-PARTICLE DENSITY MATRIX: ASYMPTOTICS AND NUMERICS

In this Section we analyze the short and long-distance asymptotics of the determinant representations of the density matrix (10). We also describe the numerical procedure which deals with the intermediate length scales. The results of this Section will be used for the calculation of the momentum distribution function given in the next Section.

Since $\rho(x) = \rho(-x)$, we assume $x \geq 0$. For *short distances* ($x \ll 1$) one can easily find from Eqs. (17) and (18) that

$$\frac{\rho(x)}{\rho(0)} = 1 - \frac{x^2}{6} - \frac{\alpha}{\pi}\frac{x^3}{18} + \frac{x^4}{120} + \frac{\alpha}{\pi}\frac{11x^5}{2700} + \dots. \tag{21}$$

For $\alpha = 0$ this is simply the expansion of $\sin x/x$. Notice that higher order terms that are not exhibited on the r.h.s. of Eq. (21), depend on α nonlinearly. Next, consider the *long-distance* expansion of Eq. (17). Compared to the short-distance analysis, this is a more sophisticated task. For the boson gas at $T \ll T_0$ the analysis carried out in Refs. [21] shows the power-law decay of the density matrix:

$$\frac{\rho(x)}{\rho(0)} = \frac{C}{\sqrt{x}}\left[1 + \frac{1}{8x^2}\left(\cos 2x - \frac{1}{4}\right) + \frac{3}{16x^3}\sin 2x\right] \tag{22}$$

with relative corrections of the order of x^{-1}. The constant C is given by $C = \pi e^{1/2} 2^{-1/3} A^{-6}$ with $A = 1.2824271\dots$ being Glaisher's constant. For the two-component fermion gas at $T \gg T_0$ the decay is exponential [20]:

$$\rho(x) = C_1 e^{-2\nu x} x^{\Delta_{\mathrm{F}}} \sin(2x - \nu \ln x + C_2). \tag{23}$$

Here C_1 and C_2 are some constants which are given explicitly in Ref. [22], while

$$\Delta_{\mathrm{F}} = \frac{\nu^2}{2} - 1, \quad \nu = \frac{\ln 2}{\pi}. \tag{24}$$

The relative corrections to Eq. (23) are of the order of x^{-1}. The technique developed in the papers [15, 16] and [17] for the two-component fermion gas can be easily adopted to the case of the two-component boson gas at $T \gg T_0$. Like for fermions, the density matrix decays exponentially in the large x limit

$$\rho(x) = \text{const}\; e^{-2\nu x} x^{\Delta_{\mathrm{B}}}. \tag{25}$$

The anomalous exponent

$$\Delta_{\mathrm{B}} = \frac{1}{2}(\nu^2 - 1) \tag{26}$$

and ν is given by Eq. (24). The relative correction to Eq. (25) is of the order of x^{-1}.

Finally, we turn to intermediate values of x. To analyze the representation (17), we use instead of Eq. (18) an alternative definition of the Fredholm determinant [23]. Let V be an $L \times L$ matrix with the entries $V_{ab} = V(k_a, k_b)$. Let

$$k_a = \left(\frac{2a}{L-1} - 1\right), \quad a = 0, 1, \dots, L-1. \tag{27}$$

Then the Fredholm determinant Eq. (18) can be represented as follows:

$$\det(\hat{I} + \hat{V}) = \lim_{L\to\infty} \det{}_L \left(I + \frac{2}{L-1}V\right). \tag{28}$$

The matrix I on the right hand side of the equation (28) is the $L \times L$ identity matrix. We use the representation (28) to calculate $\rho(x)$ for x in the range $0 \leq x \leq 20$. For the figures shown in the next Section we used the number of divisions $L = 800$.

V. LOW-TEMPERATURE RECONSTRUCTION OF THE MOMENTUM DISTRIBUTION

In this Section we present our results on the momentum distribution function of quantum two-component gases in two temperature regimes $T \ll T_0$ and $T \gg T_0$ discussed in Section II.

The momentum distribution function $n(k)$ for a one-dimensional gas is given by

$$n(k) = \int dx e^{-ikx} \rho(x), \tag{29}$$

where $\rho(x)$ is the one-particle density matrix (10). Recall that x is measured in units of $1/\sqrt{\mu}$ as defined below Eq. (7).

First, consider the gas of two-component fermions. The momentum distribution for different temperature regimes is shown in Fig. 1. The $T \ll T_0$ curve was obtained by the interpolation of the data given in Ref. [18] for the Hubbard model at low filling. The $T \gg T_0$ curve was calculated using the determinant representation discussed in Sections III and IV. One can see a dramatic change in the shape of the momentum distribution function as the temperature increases from $T \ll T_0$ to $T \gg T_0$. The singularity of the momentum distribution at $k = k_F = \pi n_{1\mathrm{D}}/2$, predicted by the Luttinger theorem



Figure 1: The momentum distribution function $n(k)$ for two-component fermions. The thick solid curve is our result for $T \gg T_0$. The dashed curve corresponds to $T \ll T_0$ and is obtained by the interpolation of the data (filled circles) from Ref. [18]. The thin solid line shows the distribution function for non-interacting two-component fermions. The momentum k is normalized to the Fermi momentum $k_F = \pi n_{1D}/2$.



Figure 2: The momentum distribution function $n(k)$ for two-component bosons. The solid curve is the result for $T \gg T_0$, the dashed curve is for $T \ll T_0$. For convenience we normalize the momentum k to the Fermi momentum $k_F = \pi n_{1D}/2$ of a fermion system with the same density.

[24], disappears and the momentum distribution spreads out to larger k. Another very pronounced effect is that the large momentum tail of the momentum distribution gets strongly suppressed as the temperature increases. This is a counterintuitive result from the point of view of physics of weakly interacting systems, where the fraction of high energy particles grows monotonously with increasing temperature. In strongly correlated systems, however, bare particles have no overlap with elementary excitations of the system and the momentum distribution of bare particles is not directly related to the distribution of energy between the eigenmodes of the system.

Next, consider the gas of two-component bosons. Using the results of Sections III and IV we calculate the momentum distribution function Eq. (29) numerically for both $T \ll T_0$ and $T \gg T_0$. The results of the calculations are shown in Fig. 2. One can see that the momentum distribution in the ground state of bosons at $T \ll T_0$ is divergent at $k = 0$. This behavior takes place at any coupling strength γ [25, 26] and is the manifestation of the quasi-long-range order in the system, that is, quasi-condensate. The situation is different at $T \gg T_0$. Due to the exponentially decaying term in the asymptotic expression Eq. (25), the function $n(k)$ is continuous with all its derivatives for all k. The suppression of the quasi-condensate component as the temperature increases from $T \ll T_0$ to above T_0 has a simple physical explanation: due to the excitation of the soft degrees of freedom the system loses coherence and cannot exhibit quasi-long-range order. For the large momentum tail of $n(k)$ one gets from Eq. (21):

$$n(k) \sim -\frac{\alpha}{k^4}, \qquad k \to \infty. \tag{30}$$

Comparing the values of α given in Table I one can see that the fraction of particles with large momentum decreases as the temperature increases from $T \ll T_0$ to above T_0 similarly to the fermionic case.

VI. DISCUSSION

One-dimensional quantum gases were recently created and studied in an optical lattice forming an array of well separated one-dimensional tubes [2]. The particle momentum distribution in such systems can be observed in experiments involving ballistic expansion [3, 4, 5]. Large values of the coupling constant, $\gamma \sim 200$, were reported in Ref. [4]. Such values of γ would give a two orders of magnitude separation between the two temperature scales T_0 and T_F, which is sufficient for the momentum reconstruction to be observable. There are, however, certain geometric limitations on the validity of the model, Eq. (1), considered in this paper, which we discuss below. For the convenience of the reader we restore the constants $\hbar$ and m in the equations.

Consider an elongated harmonic trap of transverse frequency $\omega_\perp$ and axial frequency $\omega_z = \lambda\omega_\perp$, $\lambda \ll 1$, containing a quantum gas of N particles. The characteristic length a_{1D} in Eq. (2) is related to the three-dimensional scattering length a and the transverse width $a_\perp = \sqrt{\hbar/m\omega_\perp}$ by [27]

$$a_{1D} = \frac{a_\perp^2}{a}\left(1 - C_1\frac{a}{a_\perp}\right), \tag{31}$$

where $C_1 = -\zeta(1/2)/\sqrt{2} = 1.0326\ldots$ is a numerical constant, and $\zeta(x)$ is the Riemann zeta function. Since the local chemical potential $\mu(z)$ is given by $\mu(z) = \pi^2\hbar^2 n_{1D}^2(z)/2m$, the one-dimensional density is found within the Thomas-Fermi approximation to be $n_{1D}(z) =$

$(2N - z^2/a_z^2)^{1/2}/\pi a_z$, where $a_z = \sqrt{\hbar/m\omega_z}$. The value of γ, Eq. (3), may then be expressed in terms of the particle number N, the scattering length a and the trap frequencies $\omega_\perp$ and ω_z. At the trap center ($z = 0$) one obtains the simple expression

$$\gamma(0) = \pi \left(\frac{2}{\lambda N}\right)^{\frac{1}{2}} \frac{a}{a_\perp - C_1 a}. \tag{32}$$

From Eq. (32) it is clear that to achieve large values of $\gamma(0)$ at fixed N and λ the system must be tuned to the resonance $a_\perp - C_1 a = 0$. However, very close to the resonance the scattering amplitude will no longer correspond to a delta-function potential in real space due to the presence of higher-order terms in the relative momentum. This may be demonstrated from the expansion of the scattering amplitude f as given in [27],

$$f = -\frac{1}{1 + ika_{\mathrm{1D}} - i\sqrt{2}\zeta(3/2)(ka_\perp)^3/8}. \tag{33}$$

From this expression one notes that the terms of third order in k are negligible only when $a_{\mathrm{1D}}/a_\perp \gg \sqrt{2}\zeta(3/2)(ka_\perp)^2/8$. If we identify k with the Fermi wave number k_{F} which has its maximum value $k_{\mathrm{F}} = \pi n_{1D}(0) = \sqrt{2N}/a_z$ in the center ($z = 0$) we obtain the condition

$$a_{\mathrm{1D}} \gg a_\perp \frac{\sqrt{2}\zeta(3/2)}{4} N\lambda \tag{34}$$

which in combination with Eq. (31) implies that $(a_\perp/a - C_1)/N\lambda \gg 1$, in order for our model to be applicable. Combining this result with Eq. (32) we arrive at the upper bound for the interaction constant γ for the trap with given λ and N:

$$1 \ll \gamma(0) \ll \pi\sqrt{2}\left(\frac{1}{\lambda N}\right)^{\frac{3}{2}}. \tag{35}$$

For traps with $\lambda \sim 300$ [5] containing $N = 30$ particles this estimate gives $\gamma \ll 140$.

VII. CONCLUSIONS

In conclusion, the momentum distribution of two component boson and fermion gases was considered in the limit of strong interatomic repulsion. It was shown that due to the strongly correlated nature of the system the presence of the internal degrees of freedom of the atoms results in the emergence of a new low temperature decoherent state absent in one-component gases. The onset of this state with increasing temperature is marked by a pronounced change of the momentum distribution of the atoms.

Acknowledgement

The authors would like to thank J. Dalibard for helpful discussions. M.B. Zvonarev's work was supported by the Danish Technical Research Council via the Framework Programme on Superconductivity.

[1] A. Görlitz *et al.*, Phys. Rev. Lett. **87**, 130402 (2001); F. Schreck, *ibid.* **87**, 080403 (2001); S. Dettmer, *ibid.* **87**, 160406 (2001);

[2] H. Moritz, T. Stöferle, M. Köhl, and T. Esslinger, Phys. Rev. Lett. **91**, 250402 (2003); B. Laburthe Tolra *et al.*, Phys. Rev. Lett. **92**, 190401 (2004).

[3] S. Richard, Phys. Rev. Lett. **91**, 010405 (2003).

[4] B. Paredes *et al.*, Nature **429**, 277 (2004).

[5] M. Köhl *et al.*, e-print: cond-mat/0406397.

[6] V.E. Korepin, N.M. Bogoliubov and A.G. Izergin, *Quantum Inverse Scattering Method and Correlation Functions* (Cambridge University Press, Cambridge, England, 1993).

[7] E.H. Lieb and W. Liniger, Phys. Rev. **130**, 1605 (1963); E.H. Lieb. *ibid.* **130**, 1616 (1963).

[8] L. Tonks, Phys. Rev. **50**, 955 (1936); M. Girardeau, J. Math. Phys. (N.Y.) **1**, 516 (1960).

[9] A.G. Izergin and A.G. Pronko, Nucl. Phys. B **520**, 594 (1998).

[10] C.N. Yang, Phys. Rev. Lett. **19**, 1312 (1967).

[11] E. Lieb and D. Mattis, Phys. Rev. **125**, 164 (1962).

[12] You-Quan Li *et al.*, Europhys. Lett. **61**, 368 (2003).

[13] M.A. Cazalilla and A.F. Ho, Phys. Rev. Lett. **91**, 150403 (2003).

[14] A.O. Gogolin, A.A. Nersesyan, and A.M. Tsvelik, *Bosonization and Strongly Correlated Systems* (Cambridge University Press, Cambridge, England, 1998).

[15] Vadim V. Cheianov and M.B. Zvonarev, Phys. Rev. Lett. **92**, 176401 (2004).

[16] Vadim V. Cheianov and M.B. Zvonarev, J. Phys. A: Math. Gen. **37**, 2261 (2004).

[17] G.A. Fiete and L. Balents, e-print: cond-mat/0403744.

[18] M. Ogata and H. Shiba, Phys. Rev. B **41** 2326 (1990).

[19] A. Lenard, J. Math. Phys. (N.Y.) **5**, 930 (1964).

[20] A. Berkovich and J.H. Lowenstein, Nucl. Phys. B **285**, 70 (1987).

[21] K.B. Efetov and A.I. Larkin, Sov. Phys. JETP **42**, 390 (1976); H. G. Vaidya and C. A. Tracy, Phys. Rev. Lett. **42**, 3 (1979); M. Jimbo, T. Miwa, Y. Mori, and M. Sato, Physica (Amsterdam) **1D**, 80 (1980).

[22] A. Berkovich, J. Phys. A **24**, 1543 (1991).

[23] V.I. Smirnov, *A course of higher mathematics*, Vol IV, p. 24 (Pergamon, Oxford, 1964).

[24] J.M. Luttinger, Phys. Rev. **119**, 1153 (1960); K.B. Blagoev and K.S. Bedell, Phys. Rev. Lett. **79**, 1106 (1997).

[25] V.N. Popov, JETP Lett, **31**, 526 (1980).

[26] F.D.M. Haldane, Phys. Rev. Lett. **47**, 1840 (1981).

[27] M. Olshanii, Phys. Rev. Lett. **81**, 938 (1998).